

Dantrolene Sodium

【IIDAN】 Dantrolen IV® 20mg/Vial

ATC Code : M03CA01

中文名：蘭特益靜脈注射液 «富洋»

適應症：治療惡性高熱症。

藥理分類：Skeletal Muscle Relaxant.

作用機轉：Acts directly on skeletal muscle by interfering with release of calcium ion from the sarcoplasmic reticulum; prevents or reduces the increase in myoplasmic calcium ion concentration that activates the acute catabolic processes associated with malignant hyperthermia.

用法用量：**Administration:** administered by continuous rapid intravenous push.

Dosage regimen:

Malignant Hyperthermia reaction (adults & children):

— 1 mg/kg IV by continuous rapid intravenous push and continue until symptoms subside or to a max cumulative dose of 10 mg/kg; discontinue all anesthetic agents and administer 100% oxygen (recommended).

— Prophylactic dose:

2.5 mg/kg IV, starting approximately 1.25 hr before anesthesia and infused over 1 hr.

不良反應：暈眩、困倦、發疹、腹瀉、嘔吐、肌無力。

注意事項：1.應以所附稀釋劑配製。藥品溶解後需過濾後才可使用。
2.配製後於 20-25°C 下避光貯存，可保存 6 小時；
3.每 vial 限以 60 ml 無菌注射用水配製，需充分搖震直到為澄清溶液。不可使用 D5W 及 NS。

懷 孕 期：1. Adverse events have been observed in animal reproduction studies. Dantrolene crosses the human placenta.
2. Adverse events have been observed in animal reproduction studies. Dantrolene crosses the human placenta.

授 乳 期：1. Low amounts of dantrolene are present in breast milk.
2. Due to the potential for serious adverse reactions in the nursing infant, a decision be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of treatment to the mother.

安 定 性：配製後室溫避光 6 小時內使用完畢。

儲 存：未開封請儲存於室溫(20~25°C)。避光儲存。

備 註：1.Dantrolene 列屬孤兒藥(orphan drug)，高屏地區庫存醫院有高醫、高榮、高雄長庚醫院、小港、大同、義大等。(資料來源：台灣麻醉醫學會)
2.惡性高熱諮詢專線(07)731-7123 轉 5041—高雄長庚醫院麻醉部。(麻醉科 24 小時不關機手機：0975056933) (資料來源：台灣麻醉醫學會)
3.本院現有備品係由屏東基督教醫院逕自國外採購，分售本院使用；本品僅備於開刀房使用，本院門(急)診藥局、住院藥局及藥庫皆不備其庫存。