

Calcium Chloride

【ICAL】 Vitacal[®] 20mL/Amp

ATC Code : B05BB02

中文名：美達加祿注射液 «信東»

適應症：止血、營養。

成分： Each mL contains:

Glucose · H₂O 100 mg
Calcium Chloride · 2H₂O 20 mg
(Equivalent to Ca²⁺ 0.272 mEq/mL)

藥理分類：**Calcium Salt; Electrolyte Supplement, Parenteral.**

用法用量： Adults: 10-20 mL slow IV, QD or QOD.

不良反應：若注射速度過快易發生熱潮紅、味覺異常。

交互作用：

- Calcium Chloride may ↑ the adverse/toxic effect of **Calcium Acetate**.
- Calcium Salts (Intravenous) may ↑ the adverse/toxic effect of CefTRIAxone. Ceftriaxone binds to calcium forming an insoluble precipitate.
- Calcium Salts may ↓ the therapeutic effect of Calcium Channel Blockers.

注意事項：須緩慢靜脈注射。

懷孕期：1. Calcium crosses the placenta.
2. ...antidotes should be administered to pregnant women if there is a clear indication for use and should not be withheld because of fears of teratogenicity (Bailey 2003).

授乳期：1. Calcium is excreted in breast milk.
2. The amount of calcium in breast milk is homeostatically regulated and not altered by maternal calcium intake. Calcium requirements are the same in lactating and nonlactating females (IOM 2011).

輸液相容性： Calcium chloride 與 Fat emulsion 10% , IV 不相容。

儲存：於 15~30°C 下避光貯存。勿冷凍。