

Diphenidol

【ICEP】Kphadol® 22.4mg/1mL/Amp

ATC Code : N07CA91

中文名：凱服樂注射液 «瑞士»

適應症：因內耳障礙引起之眩暈（包括腦血管障礙及頭頸部外傷後遺症之眩暈）。

【OCEP】Cephadol (Diphenidol®) 25mg/Tab

ATC Code : N07CA91

中文名：敵芬尼朵糖衣錠 «生達化學»

適應症：內耳障礙引起之眩暈（包括血管障礙及頭頸部外傷後遺症之眩暈）、（美尼攸氏）症候群。

藥理分類：Antiemetic.

用法用量：**Parenteral:**

—IM, initial 1-2 mL, give another 1 mL after 1 hr as needed, then 1-2 mL Q4H as needed.

—IV, initial 1 mL, give 1 mL after 1 hr as needed.

Tablets: orally taken after meals.

1-2 tablets TID.

不良反應：口乾，胃腸不適，視線模糊，嗜睡，頭暈，睡眠障礙，幻覺，意識混亂等。

注意事項：

1. 禁用於無尿症及腎功能障礙之患者。
2. 禁用於孕婦及6個月(或25磅)以下之嬰兒。
3. 肌肉注射:肌肉深部注射。靜脈注射:徐徐靜脈內注射。
4. 不適於作皮下注射；有過心竇性心博過速之病患，不可靜脈注射投與。

懷 孕 期：

1. Little data is available on the use of diphenidol in pregnant women.
2. Until more information is available, diphenidol should only be used during pregnancy if the maternal condition justifies the potential risk to the fetus. (Micromedex, 2020)

授 乳 期：

1. No reports describing the use of diphenidol during human lactation are available.
2. Until more data are available, use caution when considering the use of diphenidol in lactating women. (Micromedex, 2020)