

Dextran 40; Low Molecular Dextran Dextrose

【IDEX】Low Molecular Dextran Dextrose® 10% 500mL/Bot

ATC Code : B05AA

中文名：低分子血賜多朗滴注射液 «臺灣大塚»

適應症：出血時之血漿增量劑。

成分：Each 100 mL contains:

Dextran 40 10 g

Dextrose 5 g

藥理分類：**Plasma Volume Expander, Colloid.**

用法用量：**Shock:**

IV infusion, do not exceed 20 mL/kg (2 g/kg) during the first 24 hours, then $\leq$ 10 mL/kg/day. Therapy should not continue for more than 5 days. Dehydration should preferably be corrected before Dextran 40 is administered.

Treatment of thromboembolic disorders:

500 to 1000 mL over 4-6 hours on the first day, then 500 mL over 4-6 hours on the next and subsequent alternate days for not more than 10 days.

Prophylaxis of postoperative thromboembolic disorders:

500 mL over 4-6 hours may be given during or at the end of surgery and the dose repeated on the next day; treatment may be continued in high risk patients on alternate days for up to 10 days.

不良反應：噁心、嘔吐、急性腎衰竭、出血、血小板減少症、嚴重過敏反應。

注意事項：1.靜脈注射給藥。2.開栓使用後，仍有殘餘不可再使用。

懷 孕 期：Animal reproduction studies have not been conducted.

授 乳 期：It is not known if dextran is excreted in breast milk.

The manufacturer recommends that caution be exercised when administering dextran to nursing women.

儲 存：於 30°C 以下室溫，溫度變化少的環靜下儲存。勿置於冷氣口，建議以瓶子橫躺方式保存。