

Urokinase 60000U

【IURO】Urokinase-Green Cross® 60000 IU/Vial

ATC Code : B01AD04

中文名：血栓溶素注射劑 «綠十字»

適應症：急性冠狀動脈栓塞、清潔靜脈導管、急性肺栓塞、末梢動脈靜脈栓塞症。

藥理分類：**Thrombolytic agents.**

用法用量：IV or intracoronary infusion only.

Coronary artery thrombosis:

After the administration of rapid IV heparin 2500-10000 units, the prepared urokinase solution should be infused into the thrombosed coronary artery at a rate of 6000 IU/min for up to 2 hours.

Occluded IV catheters:

Gently inject 1mL of urokinase (5000 IU/mL) solution by a 1 mL tuberculin syringe; aspirate contents after 5 min and may be repeated every 5 min until patency is restored.

Pulmonary embolism:

Loading dose, 4400 IU/kg IV infusion over 10 min, followed by continuous infusion of 4400 IU/kg/hr for 12 hours.

Peripheral arterial-venous occlusion:

60,000-240,000 IU on first day of treatment followed by tapered dose for about 7 days.

Safety and efficacy not established in children.

不良反應：出血傾向、休克、S-GOT 與 S-GPT 值升高、噁心、嘔吐、厭食、發熱、惡寒、頭痛等。

交互作用：

- Agents with Antiplatelet Properties may ↑ the anticoagulant effect of Urokinase.
- Urokinase may ↑ the anticoagulant effect of Anticoagulants.
- **Tranexamic Acid** may ↓ the therapeutic effect of Thrombolytic Agents.
Thrombolytic Agents may ↓ the therapeutic effect of Tranexamic Acid.
- Urokinase may ↑ the adverse/toxic effect of **ACE Inhibitors**, specifically, the risk of angioedema.
- **Herbal Products** with Anticoagulant/Antiplatelet Effects (Bilberry, Bromelain, Emblica, Garlic, Ginger, Ginkgo Biloba, Ginseng (American), Ginseng (Panax), Ginseng (Siberian), Policosanol: ↑ the adverse effect of Thrombolytic Agents

注意事項：本品溶解後須立即使用。

懷 孕 期：可能導致胎盤未成熟前分離。唯有在效益高於風險時才可使用。

授 乳 期：尚未知 urokinase 是否會分泌至乳汁中。

相容輸注液：D5W、NS。