Lidocaine HCl

【IXYL25】Xylocaine（Lignocaine®）2% for Intravenous Injection 100mg/5mL/Amp

ATC Code：C01BB01

中文名： 里格卡因注射液 2%　«Fresenius Kabi»

適應症： 心室性不整律之急性治療。

用法用量： Administer IV bolus dose at a rate of 25-50 mg/min (0.35-0.7 mg/kg/min).

Ventricular arrhythmias:

－Adults:

Initially, 50-100 mg (0.7-1.4 mg/kg) as bolus IV injecttion. If desired response is not achieved, 25-50 mg may be administered 5 minutes after completion of the first injection, or 0.5-0.75 mg/kg may be repeated every 5-10 minutes (or every 3-5 minutes in the treatment of refractory VT or VF) as necessary, up to a maximum total loading dose of 3 mg/kg. Maximum total dose of 200-300 mg over a 1-hour period.

Maintenance infusion of 20-50 mcg/kg/min, MAX 4 mg/min.

－Children:

Initially, 0.5-1 mg/kg as bolus IV injection; dose may be repeated according to patient response, up to a maximum total dose of 3-5 mg/kg.

Maintenance infusion of 10-50 mcg/kg/min.

不良反應： 嗜睡、眩暈、焦慮、激動、視力模糊、噁心、嘔吐等。

授 乳 期： 限於判斷其治療上的有益性超越危險性時才可給藥。

相容輸注液： D5W、D5NS、D5W-1/2NS、NS、1/2NS、Lactated Ringer's、Dextrose 5% in Lactated Ringer's。