

Ritodrine HCl

【IYUT】Anpo® 50mg/5mL/Amp

ATC Code : G02CA01

中文名：安寶注射液 «信東»

適應症：預防早產、流產。

藥理分類：Beta-2 agonist, Inhibition of acute preterm labor agent.

用法用量：Administration : IV, IM

Dosage regimens :

IV infusion

- Initially 50 mcg/min, to be gradually increased by 50 mcg/min every 10 minutes until the desired response is attained.
- The effective dose is usually between 150 to 350 mcg/min. The infusion should be continued for 12 to 24 hrs after uterine contractions cease.

IM injection

- 10 mg every 3 to 6 hrs for 24 hours, with transition to oral therapy after the last IM injection.
- IM regimens avoid large infusion volumes, and therefore may have contributed to less disruption in sodium, potassium, and fluid imbalance. Maternal tachycardia is less with IM use.

不良反應：胸痛，呼吸困難，或視覺模糊，頭昏，口乾，皮膚乾、潮紅，心悸，頻尿，嘔吐，低血鉀。

注意事項：

1. 稀釋用靜脈輸注液通常為 **D5W**，糖尿病患者改用 NS。
2. 用藥期間須監測血壓及血鉀。

懷 孕 期：

1. Ritodrine can cross the placenta.
2. There are no well-controlled studies using ritodrine before the 20th week of gestation, and the manufacturer recommends ritodrine not be used before that time.

授 乳 期：

1. Ritodrine is used to inhibit premature labor, and it is therefore unlikely that a lactating female would receive this drug;
2. there is insufficient evidence to support or refute its safety in breastfeeding. It is not known whether ritodrine is excreted into human breast milk and the potential for adverse effects in the nursing infant from exposure to the drug are unknown.