

## Moxifloxacin HCl

【OMOSF】 Floxsafe 400 Film Coated Tablets 400mg

ATC Code : J01MA14

中文名：福樂星膜衣錠 400 毫克

適應症：用於治療成人(18 歲以上)感受性細菌引起的感染症，包括：上呼吸道及下呼吸道感染(急性鼻竇炎、慢性支氣管的急性惡化、社區性肺炎)，皮膚和軟組織的感染，複雜腹腔內感染(包括多種細菌感染症)。

藥理分類：**Antibiotic, Fluoroquinolone; Antibiotic, Respiratory Fluoroquinolone.**

用法用量：**Administration:**

Administer orally **without regard to meals.**

### Indications and Dosage in Adults:

**NOTE:** No dosage adjustment needed when switching from IV to oral administration, or vice versa.

#### Respiratory Tract Infections:

- Acute Sinusitis: 400 mg QD for 10 days.
- Acute Bacterial Exacerbations of Chronic Bronchitis: 400 mg QD for 5 days.
- Community-acquired Pneumonia (CAP): 400 mg QD for 7-14 days.

#### Skin and Skin Structure Infections:

- Uncomplicated Infections: 400 mg QD for 7 days.

#### Complicated intra-abdominal infections (including polymicrobial infections):

400 mg QD for 5-14 days; in geriatric patients, therapy should be initiated with IV formulation.

#### Tuberculosis:

##### Drug-susceptible tuberculosis (alternative agent):

**400 mg QD** in combination with additional appropriate antituberculosis agents.

##### Drug-resistant tuberculosis:

- **400 mg QD** in combination with additional appropriate antituberculosis agents;
- **doses of 600 or 800 mg QD** have been used in select cases (eg, elevated minimum inhibitory concentration or malabsorption).
- **Duration:** Individualize based on rapidity of culture conversion, extent of disease, and patient-specific factors, including clinical response and toxicity.

**Dosage adjustments** in Renal Impairment / Mild or Moderate Hepatic Impairment: NOT required.

**Dosage adjustments** in geriatrics: NOT required.

Safety and efficacy not established in children < 18 yr.

不良反應：胃腸不適、腹瀉、眩暈、頭痛、倦怠感、過敏反應。

交互作用：

- DRONEDARONE, DONEPEZIL, HYDROXYCHLOROQUINE, VORICONAZOLE, METRONIDAZOLE, SERTRALINE, QUETIAPINE, SULPIRIDE, FLUOXETINE, BUPRENORPHINE, DOMPERIDONE, HYDROXYZINE: ↑ risk of QT-interval prolongation.
- DEXAMETHASONE, HYDROCORTISONE, METHYLPREDNISOLONE, TRIAMCINOLONE, CORTISONE: ↑ risk of tendon rupture.
- WARFARIN: ↑ risk of bleeding.
- POTASSIUM SULFATE/MAGNESIUM SULFATE/SODIUM SULFATE: ↓ absorption of fluoroquinolone antibiotics.
- ALUMINUM, CALCIUM OR MAGNESIUM CONTAINING PRODUCTS: ↓ oral fluoroquinolone effectiveness.
- ANTIDIABETIC AGENTS: ↑ risk of hypoglycemia or hyperglycemia.
- RIFAMPIN: ↓ moxifloxacin exposure and plasma concentrations.

- ANTACIDS, SUCRALFATE, IRON: ↓ oral moxifloxacin effectiveness.

**儲存：** 應於 25°C 以下、乾燥處貯存。

**懷孕期：** 禁用。

**授乳期：** 禁用。

**健保規定：** 限用於成人(18 歲以上)之 1.慢性支氣管炎的急性惡化、社區性肺炎、複雜性腹腔內感染。2.急性鼻竇炎或複雜性皮膚和軟組織感染，因抗藥性細菌感染及病情需要，經感染科醫師會診確認需要使用者。3.若使用於結核病不予給付。