

Fenoterol HBr

【OBEA】Asmatin® 2.5mg/Tab

ATC Code : R03CC04

中文名：喘必定錠 «杏輝»

【OFENL】Strolin® Liquid 0.5mg/mL, 60mL/Bot

ATC Code : R03CC04

中文名：適喘寧液 «晟德»

適應症：下列支氣管痙攣疾患之預防與治療：支氣管氣喘、阻塞性支氣管炎、慢性支氣管炎、氣腫和伴有支氣管痙攣之肺支氣管障礙。

藥理分類：Beta2 Agonist.

用法用量：Adults, >14yrs adolescent:

Tablet: 2.5-5 mg 3 times daily before meals

Syrup: 5-10 mL TID

Children:

Tablet:

6-14 yrs, 2.5 mg 3 times daily before meals

1-6 yrs, 1.25-2.5 mg 3 times daily before meals

Syrup:

6-14 yrs: 5 mL TID

1-6 yrs: 2.5-5 mL TID

<1 yr: 2.5mL BID

不良反應：震顫、不安、心悸、眩暈、疲勞、頭痛、出汗、口乾等。

注意事項：1.建議飯前服用。
2.本品對子宮收縮具有抑制作用，故孕婦須謹慎用藥。
3.糖尿病患者須小心使用。

禁忌：甲狀腺機能亢進、瓣膜下主動脈狹窄、心跳過速之心律不整之患者禁用本劑。

懷孕期：1. Bronchodilators are regarded as relatively safe for use during pregnancy.
2. The general consensus view is that the management of asthma in pregnant patients should be the same as that of nonpregnant patients, and that poor oxygenation due to uncontrolled asthma represents a greater danger to the fetus than any potential harm associated with the drugs used to treat the disease (Weinberger & Weiss, 1995).

授乳期：1. It is not known whether fenoterol is excreted into human breast milk and the potential for adverse effects in the nursing infant from exposure to the drug are unknown.
2. It is not known if fenoterol affects the quantity or composition of breastmilk. Until more data is available, use caution when considering the use of fenoterol in lactating women.