Blopress Plus

【OBLOP】 Blopress® Plus 16/12.5 Tablet *ATC Code：C09DA06*

中 文 名： 博脈舒加強錠 《台灣武田》

適 應 症： 本態性高血壓（以 Candesartan cilexetil 或 Hydrochlorothiazide 單獨治療無法有效控制的本態性高血壓）。

成 分： Each tab contains:

Candesartan Cilexetil ..... 16 mg

Hydrochlorothiazide ..... 12.5 mg

藥理分類： Angiotensin II Receptor Blocker, diuretic ; Antihypertensive agent.

用法用量： **Administration:** orally, administer with or without food.

Dosage regimen:

One tablet once daily.

Dose of candesartan cilexetil should be titrated before switching to 16/12.5 mg.

不良反應： 背痛、上呼吸道感染、頭痛、噁心、暈眩、頭痛、高尿酸。

交互作用： **Candisartan---**

1. **Aliskiren**: ↑risk of hyperkalemia, renal impairment, and hypotension.
2. **ACEIs**: ↑risk of adverse events (ie, hypotension, syncope, hyperkalemia, changes in renal function, acute renal failure).
3. **Digoxin**: ↑risk of digoxin toxicity (nausea, vomiting, arrhythmias).
4. **Cellcept®**: ↓ mycophenolate exposure and possible reduced efficacy.
5. **Lithium**: ↑risk of lithium toxicity.
6. **Trimethoprim**, **potassium** citrate: ↑risk of hyperkalemia.

Hydrochlorothiazide—

1. **DIGITALIS** GLYCOSIDES: may result in digitalis toxicity (nausea, vomiting, arrhythmias).
2. **METHOTREXATE**: ↑methotrexate exposure and enhanced myelosuppression.
3. **TOPIRAMATE**: ↑ topiramate exposure.
4. **NSAIDS**: ↓ diuretic effectiveness and possible nephrotoxicity.

注意事項：

1. 飯前或飯後服用均可。
2. 若有經常咳嗽，應告知醫師。
3. 不建議使用於肝功能受損或嚴重腎功能受損（CCr<30 mL/min）之病患。
4. 對 **sulfonamide** 衍生物過敏者禁用本劑。
5. 痛風患者不建議使用。

懷 孕 期：

1.不建議使用於初期妊娠。禁用於懷孕第二期及第三期。

2. **[US Boxed Warning]: Drugs that act on the renin-angiotensin system can cause injury and death to the developing fetus. When pregnancy is detected, discontinue as soon as possible.**

授 乳 期：

1.由於缺乏哺乳期間使用 Blopress® Plus 的資訊，故哺乳時不建議使用 Blopress® Plus，尤其是為新生兒或早產兒哺乳期間。

2. Candesartan is present in breast milk.

3. Due to the potential for serious adverse reactions in the breastfeeding infant, the manufacturer recommends a decision be made whether to discontinue breastfeeding or to discontinue the drug, considering the importance of treatment to the mother.