Cal. Gluconate, Cal. Phosphate, Vit. D<sub>2</sub>

[OCALG] Calcium Gluconated Tab ATC Code: A12AX

中文名:葡萄糖鈣丁片錠 《昇銘藥品》

適應症: 鈣、磷質缺乏引起之軟骨症、皮膚病、妊婦、哺乳期鈣質之補給。

主成分: Each mL contains:

藥理分類: Calcium Salt; Electrolyte Supplement, Oral; vitamin d, fat soluble.

用法用量: Note: 1 Tab contains elemental Ca: 131.34 mg

**Administration:** orally, taken with plenty of fluids with or following meals.

Indications and dosage regimen:

Adults: 1 Tab three times a day with meals.

不良反應:高血鈣症、便秘。腸胃不適。

交互作用:

- Concurrent use with DIGITOXIN may result in cardiotoxicity.
- Concurrent use with Sod. POLYSTYRENE Sulfonate (eg. Kalimate) may result in an **increased** risk of systemic alkalosis.

注意事項:作為鈣質補充時,建議隨餐與食物併服。

- 懷 孕 期: 1.Calcium crosses the placenta. The ergocalciferol (vitamin D<sub>2</sub>) metabolite, 25(OH)D, crosses the placenta.
  - 2. The maternal dose of vitamin D needed to provide the infant with an adequate amount of vitamin D is still under study (*Wagner 2008*).
- 授 乳 期: 1.Calcium is present in breast milk. The 25(OH)D metabolite can be detected in breast milk
  - 2. Calcium requirements are the same in lactating and nonlactating females (*IOM*, 2011).
  - 3. Vitamin D supplementation is recommended in all infants who are partially or exclusively breast fed (*Misra 2008; Wagner 2008*).

備 註:1、鈣含量:

Calcium Gluconate......194mg\*9%=17.46 mg Ca Calcium Phosphate Dibasic... 292mg\*39%=113.88 mg Ca 1 Tab contains Ca: 17.46 mg+113.88 mg=131.34 mg

2、Vitamin D<sub>2</sub> 需求: 400-1000 IU/Day