

Activated Charcoal

【OCARB】 Carbomix(活性炭) (Carbomix Granules for Suspension®) 61.50 GM

ATC Code : A07BA01

中文名： 蓋抹滅顆粒劑 «科懋»

適應症： 緊急治療藥物或化學品中毒之病人。

用法用量： Extemporaneously, mix powder with sufficient water (e.g., 20-30 g in at least 240 mL) to form a slurry.

Poisonings:

— Adults & Adolescents \geq 13 yr:

Single dose: 25-100 g ORALLY.

Multiple dose:

50-100 g ORALLY, then 12.5 g every hour, 25 g every 2 hrs, or 50 g every 4 hrs. Continue multiple-dose therapy until patient recovers or major toxicity resolves.

— Pediatric Patients:

Single-Dose or Multiple-Dose Regimens for Treatment of Poisoning in Children

Age	Single Dose	Multiple Dose
Infants up to 1 yr of age	10-25 g or 0.5-1 g/kg	
Children up to 13 yrs of age	25-50 g or 0.5-1 g/kg	10-25 g initially, then 1-2 g/kg Q2-4h

GI Disorders:

Adults, 0.6-5 g ORALLY as a single dose or 0.975-3.9 g ORALLY 3 times daily after meals.

不良反應： 嘔吐、腹瀉、便秘、解黑便。

注意事項： 本劑對腐蝕性物質引起之中毒無效。

授乳期： 授乳婦使用本劑對被哺乳的嬰兒不會造成任何影響。

**全民健保藥品給付規定：Carbomix®限藥物中毒急救使用。