## Castor oil

[OCAST] Castor oil (Oleum Ricini®) 30mL/Bot ATC Code: A06AB05

中文名: 蓖麻子油 《應元》

適應症: 瀉劑。

藥理分類: Laxative, Miscellaneous

用法用量: Administration:

Administer orally.

Do not administer at bedtime because of rapid onset of action.

 Should be administered on an empty stomach with juice or carbonated beverages.

## **Indications and dosage regimens:**

## **Colonic evacuation:**

Administer as a single dose about 16 hours before the surgery or procedure.

-Adults: 15-60 mL

—Children: < 2 yrs, 1-5 mL; 2-11 yrs, 5-15 mL;

≥ 12 yrs, 15-60 mL

## **Constipation:**

- Adults: 15 mL, but rarely indicated for occasional constipation.

不良反應: 腹痛、噁心、嘔吐。

注意事項: 1.阻塞性腸炎、闌尾炎、腹膜炎禁用。

2. 若於睡前使用,應預留排便時間。(本品藥效作用快)

3.不可連續使用一週以上。

4. 搭配足量開水。若覺口感不佳,可併服果汁或碳酸飲料。

懷 孕 期: 孕婦不宜使用本品。

Ingestion of castor oil may be associated with induction of labor.

Use of castor oil as a laxative during pregnancy should be avoided (Cullen, 2007;

Hall, 2011; Wald, 2003).

授 乳 期: 尚未知 castor oil 是否會分泌至母乳中。