

Castor oil

【OCAST】Oleum Ricini® 30mL/Bot

ATC Code : A06AB05

中文名：蓖麻子油 «應元»

適應症：瀉劑。

藥理分類：**Laxative, Miscellaneous**

用法用量：Administer orally. Do not administer at bedtime because of rapid onset of action. Should be administered on an empty stomach with juice or carbonated beverages.

**Colonic evacuation:**

Administer as a single dose about 16 hours before the surgery or procedure.

—Adults: 15-60 mL

—Children: < 2 yrs, 1-5 mL; 2-11 yrs, 5-15 mL;

≥ 12 yrs, 15-60 mL

**Constipation:**

—Adults: 15 mL, but rarely indicated for occasional constipation.

不良反應：腹痛、噁心、嘔吐。

注意事項：  
1.阻塞性腸炎、闌尾炎、腹膜炎禁用。  
2.若於睡前使用，應預留排便時間。(本品藥效作用快)  
3.不可連續使用一週以上。  
4.搭配足量開水。若覺口感不佳，可併服果汁或碳酸飲料。

懷 孕 期：孕婦不宜使用本品。

Ingestion of castor oil may be associated with induction of labor.

Use of castor oil as a laxative during pregnancy **should be avoided** (*Cullen, 2007; Hall, 2011; Wald, 2003*).

授 乳 期：尚未知 castor oil 是否會分泌至母乳中。