Celecoxib

COCEL2 Celebrex[®] Capsule 200mg

中文名: 希樂葆膠囊 «Pfizer»

(OCELE) Celecoxib Sandoz Capsule 200mg ATC Code: M01AH01

中文名: 賽樂治山德士膠囊 200 毫克 《Sandoz》

適應症: 緩解骨關節炎之症狀與徵兆,緩解成人類風濕性關節炎之症狀與徵兆,緩解成

人急性疼痛及治療原發性經痛,緩解僵直性脊椎炎之症狀與徵兆。

藥理分類: Nonsteroidal Anti-inflammatory Drug (NSAID), COX-2 Selective.

用法用量: Administration:

• Administered without regard to meals.

 Swallowed whole or the entire contents emptied onto a teaspoon of cool or room temperature applesauce and administered immediately with water.

ATC Code: M01AH01

- The contents of the capsules sprinkled onto applesauce may be stored under refrigeration for up to 6 hours.
- Up to 200 mg BID, without regard to meals; with food, if 400 mg BID.

Indications and dosage regimens:

Osteoarthritis: 200 mg QD.

Rheumatoid arthritis in Adults: 100-200 mg BID.

Ankylosing spondylitis: 200 mg QD

- if no effect after 6 weeks, may increase to 400 mg/day.
- If no response following 6 weeks with 400 mg/day, consider discontinuation and alternative treatment.

Acute pain or Dysmenorrhea:

- 400 mg initially, followed by an additional dose of 200 mg, if needed, on the first day.
- For continued relief, 200 mg BID as needed.

Dosage adjustment:

Moderate hepatic impairment: reduce dosage by 50%.

Severe hepatic impairment: not recommended.

不良反應: 腹痛、腹瀉、消化不良、頭痛、噁心、上呼吸道感染等。

交互作用: • CYP2C9 Inducers (rifampicin, rifapentine, phenytoin, phenobarbital, carbamazepine): ↓ the serum level of Celecoxib.

 CYP2C9 Inhibitors (miconazole, amiodarone, gemfibrozil, capecitabine, fluorouracil):
 † the serum level of Celecoxib.

注意事項: ● 曾因服用 Sulfonamides、Aspirin 或其他 NSAIDs 過敏者禁用之。

● 劑量為 200 mg BID 時,不必考慮用餐時機;但是更高的劑量(400 mg BID) 應與食物一起服用,以提高吸收率。

懷孕期: 1. 於第三孕期使用 Celebrex®會增加胎兒動脈導管過早閉合的風險。

- 2. 孕婦從妊娠 30 週開始應避免使用 NSAIDs,包括 CELEBREX。
- 3. Birth defects have been observed following in utero NSAID exposure in some studies, however data is conflicting (Bloor 2013).
- 4. Use of NSAIDs can be considered for the treatment of mild rheumatoid arthritis flares in pregnant women, however use should be minimized or avoided early and late in pregnancy (Bermas 2014; Saavedra Salinas 2015).
- 授 乳 期: 1. Celecoxib 會分泌至乳汁。評估本藥對母體的重要性決定停止授乳或停藥。
 - 2. In general, NSAIDs may be used in postpartum women who wish to breastfeed; however, use should be avoided in women breastfeeding infants with platelet dysfunction or thrombocytopenia.