

Colchicine

【OCOLC】 Colchicine 0.5mg/Tab

ATC Code : M04AC01

中文名：可樂喜定錠 «鹽野義»

適應症：痛風。

藥理分類：Antigout Agent.

用法用量：Administraion : orally, taken with or without food.

Indications and dosage regimens :

Gout, treatment (acute flares):

3~4mg daily divided in 6-8 times .

Gout, prophylaxis during initiation of urate-lowering therapy:

0.5 mg QD or BID (maximum: 1 mg/day).

In patients with GI intolerance, 0.5 mg every other day.

Pericarditis, acute:

≥ 70 kg : 1mg Q12H on *day 1*, then, 0.5 mg BID x 3 months;

<70 kg or unable to tolerate higher dosing regimens:

0.5mg Q12H on *day 1*, then, 0.5 mg QD x 3 months.

Concomitant therapy: Colchicine is adjunctive therapy to be used in combination with aspirin or an NSAID; aspirin or NSAID therapy should be continued until resolution of symptoms and then tapered gradually over several weeks.

不良反應：腹瀉、噁心、嘔吐、蕁麻疹、皮膚紅點、頭痛、肌肉無力、手指或腳趾麻木感。

交互作用：Clarithromycin, erythromycin, telithromycin: ↑colchicine level, risk of intoxication.

注意事項：1.服藥後若出現腹瀉、噁心、嘔吐、腹痛，應停藥或減量。

2.避光儲存，服用時再自鋁箔包裝剝出。

3.本品可能引起骨髓抑制之副作用，建議長期使用之患者定期檢查全血球數 (complete blood counts)。

懷 孕 期：Colchicine crosses the placenta. Colchicine may be continued during pregnancy when needed to maintain remission of rheumatic disease or treat disease flair, including patients with familial Mediterranean fever.

授 乳 期：1. The decision to continue or discontinue breastfeeding during therapy should take into account the risk of infant exposure, the benefits of breastfeeding to the infant, and benefits of treatment to the mother.
2. Reconsider use if breastfed infants develop diarrhea.
3.Colchicine is considered compatible with breastfeeding when used for the treatment of gout or familial Mediterranean fever.

儲 存：阻光儲存(光直照會變質)，服用時再自鋁箔包裝剝出。