

Danazol

【ODANA】Danol[®] 200mg/Cap

ATC Code : G03XA01

中文名：達爾諾膠囊 «衛達化學»

適應症：子宮內膜異位症、早發春機發動、囊狀慢性乳腺炎、乳腺肥大、乳痛、月經過多、纖維性囊腫性乳房疾病。

藥理分類：Androgen.

作用機轉：Suppresses pituitary output of FSH and LH, resulting in regression and atrophy of normal and ectopic endometrial tissue; decreases rate of growth of abnormal breast tissue.

用法用量：**Administration: Orally**, administer consistently with regards to food; either always administer with food or administer on an empty stomach.

Note: Initiate therapy during menstruation or ensure patient is not pregnant while on therapy.

Indications and dosage regimen:

Endometriosis:

— Moderate to severe cases:

initially, 800 mg in 2 divided doses; adjust depending on patient response.

— Mild cases:

initially, 200-400 mg given in 2 divided doses; adjust depending on patient response.

Therapy for endometriosis must continue uninterrupted for 3-6 months but may be extended for 9 months if necessary.

Fibrocystic breast disease:

100-400 mg daily given in 2 divided doses depending on patient response.

When treating fibrocystic breast disease, breast pain and tenderness are significantly relieved by the first month and eliminated in 2-3 months; elimination of nodularity usually requires 4-6 months of uninterrupted therapy.

Hereditary angioedema:

— Initial, 200 mg 2-3 times daily; individualize based on patient's clinical response.

— Maintenance, decrease initial dose by 50% or less at intervals of 1-3 months or longer; if attack occurs, daily dose may be increased by up to 200 mg.

不良反應：光敏感皮膚疹、顏面潮紅、出汗、男性化作用、陰道出血。

注意事項：1.服用本藥期間應定期監測肝功能及做血液檢查。

2.服藥期間可能月經不規則或無月經。停止服藥後 60-90 天會開始規則月經，若無，應告知醫師。

3.服用本藥後會對陽光過敏，應避免在陽光下太久，並作好防曬措施。

4.病人應謹慎注意男性化的現象。

5.每日劑量超過 400 mg 會有血清酶升高或黃疸等肝功能損害報告。

懷孕期：孕婦勿用本品。因此治療期間懷孕了，則須停藥，若繼續投與胎兒會有男性化現象，尤其女胎兒會明顯導致外生殖器陰蒂肥大和陰唇融合。

授乳期：哺乳婦勿用本品。

給付規定：全民健保藥品給付規定—Danazol 限：

1、子宮內膜異位。 2、纖維性囊腫乳房性疾患之病例使用。