

Dipyridamole

【OPERS2】Dipyridamole 25mg/Tab

ATC Code : B01AC07

中文名：待匹力達糖衣錠 «生達化學»

適應症：對於慢性狹心症之治療可能有效。

【OPERS7】Parotin® 75mg/Tab

ATC Code : B01AC07

中文名：維諾心糖衣錠 «優生»

適應症：對於慢性狹心症之治療可能有效。

藥理分類：**Antiplatelet Agent; Vasodilator.**

作用機轉：Inhibits the activity of adenosine deaminase and phosphodiesterase, which causes an accumulation of adenosine, adenine nucleotides, and cyclic AMP; these mediators then inhibit platelet aggregation and may cause vasodilation; may also stimulate release of prostacyclin or PGD₂; causes coronary vasodilation.

用法用量：**Administration:**

- For better absorbance, dipyridamole should **be taken 1 hr before or 2 hrs after meals** with a glass of water(240mL).
- Could be taken with or after meals to reduce stimulation on stomach.

Indications and dosage regimens:

Coronary artery bypass graft – Thromboembolic disorder; Prophylaxis:

100 mg QID, 2 days before surgery, then after surgery, 75 mg TID plus aspirin 325 mg TID.

Prevention of thromboembolism after cardiac valve replacement:

75-100 mg QID, as an adjunct to warfarin therapy.

慢性狹心症：25mg TID (仿單)

Safety and efficacy not established in children.

不良反應：嘔吐、腹瀉、眩暈、噁心、頭痛、肌肉痛、面紅等。

懷 孕 期：There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, Dipyridamole tablets should be used during pregnancy only if clearly needed.

授 乳 期：As dipyridamole is excreted in human milk, caution should be exercised when Dipyridamole tablets are administered to a nursing woman.