

## Oxybutynin Chloride

【ODITR】 Oxbu<sup>®</sup> Extended-Release 5mg/Tab

ATC Code : G04BD04

中文名： 歐舒緩釋錠 《健喬信元》

適應症： 治療伴有急尿、頻尿或急迫性尿失禁症狀的膀胱過動症。

藥理分類： **Antispasmodic Agent, Urinary.**

用法用量： **Administration :**

- Orally, administer without regard to meals.
- Must be swallowed whole with liquid; do not crush, divide, or chew; take at approximately the same time each day.

**Dosage regimen :**

**Adults:**

Initial: 5 to 10 mg QD; adjust dose as needed and tolerated in 5 mg increments every 1 to  $\geq 2$  weeks.

Maximum: 30 mg QD.

**Children  $\geq 6$  years and Adolescents:**

Initial: 5 mg QD; adjust dose as needed in 5 mg increments at weekly intervals.

Maximum daily dose: 20 mg

不良反應： 口渴、食慾不振、便秘、胃腸不適、頭痛、噁心、嗜睡、抑制泌乳。

禁忌： 本劑禁用於窄角性青光眼、胃腸道阻塞、泌尿道阻塞、嚴重結腸炎、重症肌無力、急性出血導致心血管情況不穩之患者。

交互作用：

- POTASSIUM: ↑ risk of gastrointestinal lesions.
- DONEPEZIL, BUPROPION: ↑ risk of seizure.
- QUETIAPINE, TIOTROPIUM: ↑ risk of anticholinergic side effects, including intestinal obstruction.
- SCOPOLAMINE: ↑ risk of CNS adverse reactions, intestinal obstruction, and urinary retention.

注意事項： 1. 飯前或飯後服用均可。  
2. 須整粒吞服，不要咀嚼或壓碎。  
3. 服藥期間盡量避免開車。

懷孕期： 1. 動物實驗顯示無畸胎之不良結果，但臨床上孕婦之安全使用報告乃未確定，所以本品須經醫師指示方可使用。

2. Adverse events were not observed in animal reproduction studies.

Information related to the use of oxybutynin in patients treated for neurogenic bladder during pregnancy is limited (*Andretta 2018*).

授乳期： 1. 授乳婦不建議給藥。服用本品可能抑制泌乳。  
2. It is not known if oxybutynin is present in breast milk. Suppression of lactation has been reported.