

Ergometrine Maleate

【OERGN】Ergonovine (Ergometrine®) 0.2mg/Tab

ATC Code : G02AB03

中文名：意如宮錠 «永信»

適應症：分娩時的子宮弛緩性出血、流產後出血、產褥期出血、子宮內膜刮除後的出血、子宮收縮不全。

藥理分類：Ergot Derivative; Oxytotic Agent.

用法用量：Administration: orally, may be taken with or without food. Usually administered after first parenteral dose.

Dosage regimen:

To minimize late postpartum hemorrhage, 0.2 to 0.4 mg every 6-12 hrs until the danger of uterine atony has passed, usually 48 hrs.

不良反應：高血壓、噁心、嘔吐、眩暈、頭痛、鼻塞等。

注意事項：避免與葡萄柚汁併服。

懷 孕 期：1.孕婦禁用。(仿單)

2. Fetal risk cannot be ruled out. Available evidence is inconclusive or is inadequate for determining fetal risk when used in pregnant women or women of childbearing potential. Weigh the potential benefits of drug treatment against potential risks before prescribing this drug during pregnancy. (MicroMedex,2020)

授 乳 期：1.哺乳期間不建議使用；若需使用則暫停哺乳。

2. Evidence and/or expert consensus has demonstrated harmful infant effects when used during breastfeeding. An alternative to this drug should be prescribed or patients should be advised to discontinue breastfeeding. (MicroMedex,2020)

儲 存：於 25°C 以下陰涼乾燥處避光儲存。