## Raloxifene HCl

[OEVIS] Evista® Film Coated Tablets 60mg/Tab ATC Code: G03XC01

中文名: 鈣穩膜衣錠 《臺灣禮來》

適應症: 預防及治療停經後婦女骨質疏鬆症。

藥理分類: Selective Estrogen Receptor Modulator (SERM).

用法用量: **Administration**: Orally, take without regard to meals.

Indications and dosage regimen:

**Osteoporosis:** 

Females: Oral: 60 mg once daily.

Invasive breast cancer risk reduction:

Female: Oral: 60 mg once daily.

Duration of therapy for breast cancer risk reduction: 5 years;

may be used longer than 5 years in females with osteoporosis where breast cancer

risk reduction is a secondary benefit (Visvanathan 2013).

不良反應: 腿部痙攣、臉潮紅、水腫、血栓。

注意事項: 1.飯前或飯後服用皆可。

2.停經前婦女不適合使用本藥。

禁 忌: 1.目前患有或曾有静脈血栓性栓塞(VTE)的婦女禁用 Evista®,包括深部静脈血

栓、肺栓塞及視網膜靜脈血栓。

2.已懷孕、可能懷孕或哺乳之婦女,禁止使用 Evista®。

懷 孕 期: 1.懷孕者或可能懷孕者請勿服用本藥。

2. Raloxifene is not indicated for use in females of reproductive potential.

3. Raloxifene is **contraindicated during pregnancy**.

授 乳 期: 1.授乳婦不應使用 Evista®。

2. It is not known if raloxifene is present in breast milk. Raloxifene is not indicated

for use in females of reproductive potential.