

Flavoxate HCl

【OGEN】 Genurin® 200mg/Tab

ATC Code : G04BD02

中文名： 尤利百舒糖衣錠 «德河»

適應症： 一般及尿道手術後或後遺之尿道痙攣、疼痛及由於導尿管或膀胱內視鏡引起之痙攣諸症之治療。

藥理分類： Antispasmodic Agent, Urinary.

用法用量： **Administration:** Orally Taken without regard to meals.

Dosage regimens:

Adults & children > 12 years:

100-200 mg 3-4 times per day; may reduce dose with improvement of symptoms.

不良反應： 頭痛、神智混亂、噁心、嘔吐、複視、思睡、心跳加速、便秘、心悸。

交互作用：

- **Ipratropium (Oral Inhalation):** May enhance the anticholinergic effect of Anticholinergic Agents. Risk X: *Avoid combination.*
- **Tiotropium:** Anticholinergic Agents may enhance the anticholinergic effect of Tiotropium. Risk X: *Avoid combination.*
- **Potassium Chloride:** Anticholinergic Agents may enhance the ulcerogenic effect of Potassium Chloride. Management: Patients on drugs with substantial anticholinergic effects should avoid using any solid oral dosage form of potassium chloride. Risk X: *Avoid combination.*
- **Potassium Citrate:** Anticholinergic Agents may enhance the ulcerogenic effect of Potassium Citrate. Risk X: *Avoid combination.*

注意事項： 1.空腹或與食物一起服用皆可。
2.幽門及十二指腸阻塞、胃腸出血、下尿道泌尿器阻塞之患者禁用本劑。
3.青光眼之患者應小心使用本劑。
4.本劑不建議使用於 12 歲以下孩童。

懷 孕 期： 1.動物試驗顯示對胎兒或孕婦無害。
2.若需服用本藥，懷孕前三個月需特別小心。
3.Adverse events have not been observed in animal reproduction studies.

授 乳 期： 1.尚不知 Flavoxate 是否會分泌至乳汁。
2.It is not known if flavoxate is excreted in breast milk. The manufacturer recommends that caution be exercised when administering flavoxate to nursing women.