

GlucoMet 5/500

【OGLUM】 GlucoMet 5/500 (GlucoMet® 5/500) Tablet

ATC Code : A10BD02

中文名： 克糖美錠 5/500 «友華»

適應症： 第二型糖尿病。

成分： Each tab contains:

Glyburide (別名 Glibenclamide) 5 mg

Metformin HCl 500 mg

藥理分類： Antidiabetic Agent, Sulfonylurea; Biguanide.

用法用量： Safety and efficacy not established in children.

Administration: orally, taken with meals; once daily with morning meal; twice daily with morning and evening meals.

Indication and dosage regimens;

Diabetes Mellitus Type 2:

— Initial therapy 1.25 mg/250 mg once or twice daily;

titrate in increments of 1.25 mg/250 mg per day every 2 weeks

— Previously treated patients:

2.5 mg/500 mg to 5 mg/500 mg twice daily; titrate in increments of 5 mg/500 mg up to MAX 20 mg/2000 mg once daily.

不良反應： 低血糖、胃腸不適、噁心、眩暈、維生素 B₁₂ 缺乏。

交互作用：

- **Leflunomide:** ↑ exposure of Glyburide.
- **Beta blockers:** hypoglycemia or hyperglycemia; ↓ symptoms of hypoglycemia
- **Fluoroquinolones:** ↑ risk of hypoglycemia or hyperglycemia.
- **Hydroxychloroquine:** hypoglycemia.
- **Iodinated contrast media:** result in lactic acidosis and acute renal failure.
- **Bupropion:** ↓ renal clearance of metformin.
- **Aspirin:** ↑ risk of hypoglycemia.
- **Clarithromycin:** ↑ plasma glyBURIDE concentrations.
- **Rifapentine:** ↓ oral hypoglycemic effectiveness.

注意事項： 1.建議隨餐服用。

2.避免大量飲酒(因酒會延長低血糖作用)。

3.因檢查要使用顯影劑時,可能需暫停服用數日,請向醫師諮詢。

懷 孕 期： 1.請勿服用本藥，否則可能會對胎兒造成傷害。建議應改用胰島素治療。

2. **Insulin** is the preferred medication for treating hyperglycemia in gestational diabetes mellitus. **Metformin** and **glyburide should not** be used as first-line agents, as both cross the placenta to the fetus. Other oral and noninsulin injectable glucose-lowering medications lack long-term safety data. (ADA 2020).

授 乳 期： 1.禁用本品。

2. Glyburide and metformin are present in breast milk.

用藥規定： metformin： (1)腎絲球體過濾率 (eGFR) 小於 30 ml/min/1.73 m² 禁用。(2)腎絲球體過濾率 (eGFR) 介於 30-45 ml/min/1.73 m² 應減量使用。(3)每日最大劑量：速效劑型為 3000 mg；緩釋劑型為 2000mg。