

Cefaclor Monohydrate

【OKEFL】Cero<sup>®</sup> 250mg/Cap

ATC Code : J01DC04

中文名： 賜福膠囊 «信東»

適應症： 甲族乙型溶血性鏈球菌、葡萄球菌、肺炎雙球菌、大腸桿菌、奇異變形桿菌、克雷氏肺炎桿菌、嗜血桿菌、淋球菌等病原菌所引起之感染。

藥理分類： Antibiotic, Cephalosporin (**Second** Generation).

用法用量：**Administration:** Orally, administer around-the-clock to promote less variation in peak and trough serum levels.

**Indications and dosage regimens:**

**Adults:**

—**Pneumonia:**

500 mg every 8 hours.

—**Acute otitis media / Pharyngitis and tonsillitis / Skin and skin structure infections / Urinary tract infections :**

250 mg Q8h. For more severe infections or those caused by less susceptible organisms, 500 mg Q8h.

**Children  $\geq$  1 month of age:**

—**Pneumonia:**

40 mg/kg/day in divided doses every 8 hours.

—**Acute otitis media:**

40 mg/kg/day in divided doses every 8 or 12 hours.

—**Pharyngitis and tonsillitis:**

20 mg/kg/day in divided doses every 8 or 12 hours for 10 days. For more severe infections or those caused by less susceptible organisms, 40mg/kg/day in divided doses every 8 hours.

—**Skin and skin structure infections / Urinary tract infections:**

20 mg/kg/day in divided doses every 8 hours. For more severe infections or those caused by less susceptible organisms, 40 mg/kg/day in divided doses every 8 hours.

Renal function impairment usually does not require a reduction in dose.

不良反應： 過敏性反應、腹瀉。

懷 孕 期： Adverse events were not observed in animal reproduction studies. An increased risk of teratogenic effects has not been observed following maternal use of cefaclor.

授 乳 期： 微量 Cefaclor 會分泌至母乳中，應小心使用。

使用規定： 全民健保藥品給付規定—Cefaclor 口服製劑用於：

1、急性化膿性中耳炎或急性化膿性鼻竇炎。2、急性下呼吸道感染。3、敗血症等適應症，得免附細菌敏感試驗報告，惟以 10 日為限。