

Potassium Gluconate

【ORAD】、k-Glu Oral Solution 20 mEq/15 mL/Vial

ATC Code : A12BA05

中文名： 鉀補口服液 《信東生技》

適應症： 鉀缺乏症。

成分： Each mL contains:
Potassium Gluconate 312mg (1.33 mEq K⁺)

藥理分類： Electrolyte Supplement, Oral.

用法用量： **Administration:** Orally, take after meals with water or juice.
15mL TID~QID

Prevention of hypokalemia during diuretic therapy:

Adult, 20-80 mEq/day in 3-4 divided doses.

Child, 1-2 mEq/kg/day in 1-2 divided doses.

Treatment of hypokalemia:

Adult, 40-100 mEq/day in 2-4 divided doses.

Child, 2-5 mEq/kg/day in 2-4 divided doses.

不良反應： 噁心、嘔吐、食慾不振、下痢、腹痛、胸悶、便秘、或高血鉀

交互作用：

- ANTICHOLINERGICS (SOLIFENACIN, BIPERIDEN, HYOSCYAMINE, TRIHEXYPHENIDYL, ATROPINE, OXYBUTYNIN): risk of gastrointestinal lesions.
- ACE inhibitors, ARBs : hyperkalemia.

注意事項：

- 1.本品為口服製劑，不得用於注射。
- 2.與開水或果汁混合後服用，飯後服用
- 3.本藥不得與會滯留鉀離子之利尿劑併用。
- 4.本藥不建議使用於孕婦及授乳婦。(仿單)
- 5.對 Addison's disease(安迪生氏病)患者，必須施以適當的安迪生氏病療法。
- 6.由於原發性 Aldosteronism(醛固酮過多症)引起之低鉀症，請與醛固酮拮抗劑併用。
- 7.腎功能障礙、腎功能低下及細尿管、輸尿管病變患者請謹慎服用。

懷 孕 期：

- 1.Adverse events have not been observed following use of potassium supplements in healthy women with normal pregnancies.
- 2.Use caution in pregnant women with other medical conditions (eg, preeclampsia; may be more likely to develop hyperkalemia) (IOM 2004).

授 乳 期：

Potassium is excreted into breast milk (IOM 2004).