

Lugol's Solution

【OLUG】Lugol's Solution 3mg(I₂)/mL

ATC Code : H03CA

成分： Iodine 0.3125 g
Potassium Iodide 0.625 g
Distilled Water qs ad 100 mL

藥理分類： Antithyroid Agent.

用法用量： Thyroidectomy:
2-5 mL TID PO for 10 days before surgery.

懷 孕 期： 1. Iodide crosses the placenta (may cause hypothyroidism and goiter in fetus/newborn).
2. Use for protection against thyroid cancer secondary to radioactive iodine exposure is considered acceptable based upon risk:benefit, keeping in mind the dose and duration. Repeat dosing should be avoided if possible.
3. Large amounts of iodine during pregnancy can cause fetal goiter or hyperthyroidism. Transient hypothyroidism in the newborn has also been reported following topical or vaginal use prior to delivery.

授 乳 期： 1. Skin rash in the nursing infant has been reported with maternal intake of potassium iodide.
2. Iodine is excreted in breast milk and is a source of iodine for the nursing infant.
3. Application of topical iodine antiseptic solutions can be absorbed in amounts which may affect levels in breast milk. Exposure to excess iodine may cause thyrotoxicosis.

儲 存： 避光冷藏。

備 註： 本品以原料 I₂ 及 KI 自行配製。