

Norethindrone Acetate

【ONORE5】Nordron® 5mg/Tab

ATC Code : G03DC02

中文名： 拿能寧糖衣錠 «井田»

適應症： 切迫性及習慣性流產、月經過多症、出血性子宮血崩、月經困難、續發性及原發性無月經、乳汁分泌之促進。

藥理分類： **Progestin.**

用法用量： **Administration** : oral, take at the same time each day.

Indications and dosage regimen:

Dysfunctional bleeding:

5 mg/day for 10 days.

Primary & secondary amenorrhea:

Estrogen first, then 5 mg BID from day 19-26 of the cycle.

Premenstrual syndrome & mastopathy:

5 mg QD-BID from day 19-26 of the cycle.

Endometriosis:

5 mg BID from day 5 of the cycle.

不良反應： 食慾不振、噁心、子宮不正常出血、下腹部痛等。

注意事項： 1.每天同一時間服用。
2.嚴重肝疾患或肝障害禁用。
3.孕婦禁用。

懷 孕 期： 1. Use is **contraindicated during pregnancy**.
2.First trimester exposure of progestins may cause genital abnormalities including hypospadias in male infants and mild virilization of external female genitalia.

授 乳 期： 1. Norethindrone is present in breast milk.
2. The manufacturer of norethindrone acetate recommends that caution be used if administered to a breastfeeding woman.