

Amlodipine Besylate

【ONORV】 Norvasc® 5mg/Tab

ATC Code : C08CA01

中文名：脈優錠 «輝瑞»

適應症：高血壓、心絞痛。

藥理分類：Antianginal Agent; Antihypertensive; Calcium Channel Blocker, Dihydropyridine.

用法用量：**Administration:** orally, administered with or without food.

Indications and dosage regimen:

Hypertension (HTN):

- Initially as monotherapy, 2.5-5 mg QD. When adding to existing antihypertensive regimen, initially 2.5 mg QD.
- Increase dosage gradually over 7-14 days until optimum control of BP is obtained. May increase more rapidly if symptoms so warrant and patient's tolerance and response are frequently assessed.
- Usual maintenance dosage is 5-10 mg QD. **Maximum** daily dose: 10mg.

Angina:

Usual dosage is 5-10 mg QD; adequate control usually required a maintenance dosage of 10 mg daily.

Special Populations:

—Hepatic Impairment:

for HTN, initially, 2.5 mg daily; for angina, initially, 5mg daily

—Geriatric Patients:

Initially, 2.5 mg daily. Adjust subsequent dosage based on patient response and tolerance; usually should **not exceed 5 mg daily**.

不良反應：頭痛、下肢水腫、頭昏、心悸、潮紅、噁心等。

注意事項：1.飯前或飯後服用均可，最好每天同一時間服用。
2.本品不可與葡萄柚(汁)併服。

交互作用：1. ↑ amlodipine 血中濃度：ketoconazole、itraconazole、ritonavir、cimetidine、clarithromycin、grapefruit(juice)。
2. ↓ amlodipine 血中濃度：Carbamazepine, phenobarbital, phenytoin, rifampicin、St. John's wort (聖約翰草)。
3. Simvastatin, Lovastatin: ↑ statins serum level and ↑ myopathy.
4. Clopidogrel: ↓ antiplatelet effect and ↑ risk of thrombotic events.

懷 孕 期：鈣離子通道阻斷劑有可能使得母體血壓過低而導致胎兒缺氧，因此除了預期效益比胎兒風險還高時，孕婦不應使用本藥物。

授 乳 期：Amlodipine 可分泌至人乳中。使用本品進行治療時應停止授乳。