

Acetaminophen, caffeine

【OPANC】Quit A.C. Analgesic Tablet

ATC Code : N02BE51

中文名：快安治痛加強錠 «壽元»

成分：Each tab contains:

Acetaminophen 500 mg

Caffeine 65 mg

藥理分類：Analgesic and Antipyretics; CNS Stimulant; Phosphodiesterase Enzyme Inhibitor, Nonselective.

用法用量：**Administration:** orally, taken without regard to food; may administer with food to decrease possible GI upset

Dosage regimens:

Adults:

1 Tab Q4~6H. Max. dose : 4 g/day

Children:

6~12 years old:

0.5 Tab Q4~6H

3~6 years old:

0.25 Tab Q4~6H

< 3 years old: consult doctor

適應症：解熱、鎮痛（頭痛、牙痛、咽喉痛、關節痛、神經痛、肌肉痛、月經痛）。

不良反應：搔癢、便秘、噁心、嘔吐。

交互作用：1. **Linezolid**: ↑ the hypertensive effect of Sympathomimetics.
2. **Probenecid**: ↑ the serum concentration of Acetaminophen.
3. **Sorafenib**: Acetaminophen may enhance the hepatotoxic effect of Sorafenib. SORafenib may increase the serum concentration of Acetaminophen.
4. May increase the serum concentration of **TiZANidine**.

注意事項：5. 飯前或飯後服用均可。
6. 兩次使用間隔請勿短於4小時。
7. 如出現黃疸、噁心、嘔吐、胃部痙攣或胃痛、上腹漲或痛等症狀應停藥並立即就醫。
8. 肝病、腎臟病、慢性酒精中毒以及老年人服用本品宜小心。
9. 服用本藥期間勿飲用含酒精物品。

懷孕期：1. Acetaminophen crosses the placenta.
2. Acetaminophen is recommended to be used at the lowest effective dose for the shortest duration of time to effectively treat the mother and protect the health of the fetus (Kilcoyne 2017).

哺乳期：1. Acetaminophen is present in breast milk.
2. Acetaminophen is one of the preferred non-narcotic agents (Sachs 2013) and is considered compatible with breastfeeding when used in usual recommended doses (WHO 2002).