

Pantoprazole

【OPANY】Pantyl® Gastro-resistant Tablet 40mg/Tab

ATC Code : A02BC02

中文名：速潰樂腸溶膜衣錠 «中化»

適應症：合併二種適當之抗微生物製劑治療與幽門螺旋桿菌相關之消化性潰瘍、胃潰瘍、十二指腸潰瘍、中度及嚴重逆流性食道炎、Zollinger-Ellison-Syndrome。

藥理分類：Proton Pump Inhibitor.

用法用量：**Administration:**

- Should be swallowed whole, do not split, crush, or chew. For most indications, administer 30 to 60 minutes before a meal (preferably); best if taken before breakfast.
- Manufacturer's labeling: tablets may be administered with or without food and concomitant administration of antacids do not affect absorption.

Dosage regimens:

Erosive esophagitis associated with GERD:

Treatment: 40 mg QD for up to 8 weeks; an additional 8 weeks may be used in patients who have not healed after an 8-week course.

Maintenance of healing: 40 mg QD (US labeling) or 20 to 40 mg QD (Canadian labeling); 20 mg once daily has been used successfully in maintenance of healing.

Note: Has not been studied beyond 12 months.

Hypersecretory disorders (including Zollinger-Ellison):

Initial: 40 mg BID;

Doses up to 240 mg daily have been administered.

不良反應：腹瀉、便秘、脹氣、頭痛。

交互作用：

- Cellcept® : ↓ exposure to mycophenolic acid (MPA), the active metabolite of mycophenolate mofetil.
- ERLOTINIB, CAPECITABINE, GEFITINIB, CILOSTAZOL: ↓ exposure of these drugs.
- METHOTREXATE : ↑ risk of methotrexate toxicity.
- CITALOPRAM: ↑ citalopram exposure and risk of QT interval prolongation.

注意事項：

1. 早上服用一次，飯前或飯後均可。
2. 不適合剝半或嚼碎，必須整顆吞服，與制酸劑合用不會影響本品的吸收。

懷孕期：Based on available data, PPIs may be used when clinically indicated.

授乳期：

1. Pantoprazole is present in breast milk (*Plante 2004*).
2. No adverse events were reported in the mother or infant for the 2 weeks following administration (*Plante 2004*).

給付規定：**全民健保藥品給付規定：依「消化性潰瘍用藥限制使用規定」辦理。