Dipyridamole

[OPERS2] Dipyridamole 25mg/Tab

中文名: 待匹力達糖衣錠 《生達化學》

適應症:對於慢性狹心症之治療可能有效。

[OPERS7] Parotin[®] FC Tab 75mg/Tab

中文名: 維諾心膜衣錠 《優生》

適應症:對於慢性狹心症之治療可能有效。

藥理分類: Antiplatelet Agent; Vasodilator.

作用機轉: Inhibits the activity of adenosine deaminase and phosphodiesterase, which causes an accumulation of adenosine, adenine nucleotides, and cyclic AMP; these mediators then inhibit platelet aggregation and may cause vasodilation; may also stimulate release of prostacyclin or PGD2; causes coronary vasodilation.

用法用量: Administration:

- For better absorbance, dipyridamole should **be taken 1 hr before or 2 hrs after meals** with a glass of water(240mL).
- Could be taken with or after meals to reduce stimulation on stomach.

Indications and dosage regimens:

Coronary artery bypass graft – Thromboembolic disorder; Prophylaxis:

100 mg QID, 2 days before surgery, then after surgery, 75 mg TID plus aspirin 325 mg TID.

Prevention of thromboembolism after cardiac valve replacement:

75-100 mg QID, as an adjunct to warfarin therapy.

慢性狹心症:25mg TID (仿單)

Safety and efficacy not established in children.

- 不良反應: 嘔吐、腹瀉、眩暈、噁心、頭痛、肌肉痛、面紅等。
- 懷 孕 期: There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, Dipyridamole tablets should be used during pregnancy only if clearly needed.
- 授 乳 期: As dipyridamole is excreted in human milk, caution should be exercised when Dipyridamole tablets are administered to a nursing woman.

ATC Code : B01AC07

ATC Code : B01AC07