#### Conjugated Estrogens

[OPREM] Premarin (Estromon®) 0.625mg/Tab ATC Code : G03CA57

中文名: 伊使蒙膜衣錠 《生達化學》

適應症: 更年期障礙、卵巢欠缺症狀、卵巢機能不全、機能性子宮出血、腔炎。

藥理分類: Estrogen Derivative.

用法用量: Administration: orally, may be taken without regard to meals.

# **Indications and dosage regimens:**

# **Estrogen replacement therapy:**

## -Vasomotor symptoms:

Initially, 0.3 mg/day continuously or in cyclic regimen (25 days on, 5 days off). Adjust dosage based on patient response.

#### -Vulvar and vaginal atrophy:

Initially, 0.3 mg/day continuously or in cyclic regimen (25 days on, 5 days off). Adjust dosage based on patient response.

Attempts to discontinue or taper medication should be made at 3- month and 6-month intervals

## Prevention of postmenopausal osteoporosis:

Initially, 0.3 mg/day continuously or in cyclic regimen (25 days on, 5 days off). Adjust dosage based on clinical and BMD response.

#### Hypoestrogenism:

- -Female castration or primary ovarian failure:
- 1.25 mg/day in a cyclic regimen (3 wk on and 1 wk off). Adjust dosage based on symptom severity and clinical response.
- -Female hypogonadism:

0.3-0.625 mg/day in a cyclic regimen (3 wk on, 1 wk off). Adjust dosage based on symptom severity and endometrial responsiveness.

# Breast cancer, for palliation only, in appropriately selected women and men with metastatic disease:

10 mg 3 times daily for at least 3 months.

**Advanced androgen-dependent carcinoma of prostate, for palliation only :** 1.25-2.5 mg three times daily.

#### **Dysfunctional uterine bleeding:**

2.5 to 5 mg/day in divided doses for 7 days, with a progestin. Conjugated estrogens 2.5 to 3.75 mg daily may be given plus Provera<sup>®</sup> 10 mg/day for 7-10 days.

不良反應: 體重改變、噁心、腹痛、陰道出血異常、乳房疼脹、白帶、下肢腫脹。

注意事項: 飯前或飯後服用均可。盡量每天同一時間服用。

懷 孕 期: 已經懷孕或可能懷孕婦女禁止使用。

授 乳 期: 藥品成分會分泌至乳汁,會影響乳汁品質及分泌量,須謹慎使用。