Protase Enteric Coated Capsules

【OPROT】Protase(Protase®)280mg/Cap ATC Code: A09AA02

中文名: 優妙化腸溶微粒膠囊 《永信》

適應症: 囊腫性纖維化疾病、慢性胰臟炎、胰臟切除、胃腸鏡道手術及因腫瘤引發胰

管式膽管阻塞等疾病所導致的胰液分泌不全。

成 分: Each capsule contains:

用法用量: Clinical experience should dictate initial starting dose.

Administration: Doses should be taken during meals or snacks, not before or after. Do not take without food.

Dosage regimens:

Adults & Children > 6 yrs:

Usual initial starting dosage is one capsule per meal or snack.

Children ≤ 6 yrs:

- —The exact dosage of Protease® Capsule should be selected based on clinical experience for this age group.
- -Cystic fibrosis:

1,500 to 3,000 lipase USP units/kg/meal. Dosage should be adjusted according to the severity of the disease, control of steatorrhea and maintenance of good nutritional status. MAX 6,000 lipase USP units/kg/meal.

禁 忌: 初期之急性胰臟炎禁用。

不良反應: 常見的有下痢、噁心、嘔吐、脹氣、便秘、胃絞痛等。過高的劑量可能導致

尿中尿酸過高及高尿酸血症。

注意事項: 1、切勿於飯前或飯後及無食物的狀態下服用。

2、服用時可將膠囊打開,但不可壓碎或咀嚼,亦不可與 pH 5.5 以上之食物混合,以免提早釋放出酵素,造成口腔黏膜刺激及失去酵素活性。

授 乳 期: 本藥是否分泌於人類乳汁中尚未確立。