

Phenazopyridine HCl

【OPYR】Urogen[®] 50mg/Tab

ATC Code : G04BX06

中文名：抑炎源糖衣錠 «強生»

適應症：膀胱炎、腎盂炎、尿道炎、泌尿生殖器粘膜之鎮痛。

藥理分類：Analgesic, Urinary.

作用機轉：An azo dye which exerts local anesthetic or analgesic action on urinary tract mucosa through an unknown mechanism

用法用量：**Administration:** administer without regard to meals.

Indications and dosage regimen:

Dysuria, for symptomatic relief of pain, burning, urgency, and frequency arising from irritation of the urinary tract (infection, trauma, surgery, endoscopic procedures, or catheter placement):

—Adults:

2 Tablets TID.

—Children 6-12 yrs:

1 Tablet Tid.

不良反應：噁心、腹痛、消化不良；頭暈、頭痛、發燒；皮膚或眼睛變黃；皮膚起疹；臉、手指、腳腫脹等。

交互作用：

- May ↑ the adverse/toxic effect of **Sodium Nitrite**. Combinations of these agents may increase the likelihood of significant methemoglobinemia.
- Nitric Oxide may ↑ the adverse/toxic effect of Methemoglobinemia Associated Agents. Combinations of these agents may increase the likelihood of significant methemoglobinemia.

注意事項：1、本品不可研磨使用。（避免沉積牙齒導致著色）
2、服用本劑後，尿液會呈黃色或黃褐色。

懷孕期：Phenazopyridine 會通過人類胎盤。動物試驗未發現不良作用。

授乳期：本品用於授乳婦時，其對嬰兒的安全性尚未確立。