Potassium Gluconate

[ORAD] Radi- $k^{\text{®}}$ 595mg/Tab (2.54 mEq of K⁺) ATC Code : A12BA05

中文名: 鉀順錠 《寶齡富錦》

適應症: 血鉀過低症(因手術前後、使用藥物、瀉肚、心臟病引起之血鉀過低症)。

成 分: Each tab contains:

藥理分類: Electrolyte Supplement, Oral.

用法用量: **Administration:** orally, taken with or after meals.

Indications and dosage regimens:

Note: If the dosage exceeds 16 mEq K⁺, it should be taken in divided doses.

Prevention of hypokalemia:

Average dosage approximately 20 mEq daily. Usually should not exceed 150 mEq daily.

mEq daily.

Treatment of hypokalemia:

Usual dosage is 40-100 mEq or more daily. Usually should not exceed 150 mEq daily.

不良反應: 噁心、嘔吐、食慾不振、下痢、腹痛、胸悶、便秘、或高血鉀

交互作用:

- Potassium-Sparing Diuretics: † the hyperkalemic effect of Potassium Salts.
- ARBs, ACEIs:
 † hyperkalemic effect of Potassium Salts.
- Heparin, low molecular weight Heparin, Nicorandil ,NSAIDs:
 † the hyperkalemic effect of Potassium Salts.

注意事項: 1.低鉀狀態時作為鉀的補充劑,使用前必須先檢查血清中鉀值、鈉值及尿量。

- 2.使用中定期檢查血清中鉀值、尿量和尿中鉀值鈉值的比率以及腎功能。
- 3.長期使用時,血中及尿中鉀值、腎功能、心電圖最好定期檢查。
- 4.對 Addison's disease(安迪生氏病)患者,必須施以適當的安迪生氏病療法。
- 5.由於原發性 Aldosteronism(醛固酮過多症)引起之低鉀症,請與醛固酮拮抗劑 併用。

6. 腎功能障擬、腎功能低下及細尿管、輸尿管病變患者請謹慎服用。

懷 孕 期: 1.Adverse events have not been observed following use of potassium supplements in healthy women with normal pregnancies.

2.Use caution in pregnant women with other medical conditions (eg, preeclampsia; may be more likely to develop hyperkalemia) (IOM 2004).

授 乳 期: Potassium is excreted into breast milk (IOM 2004).