

【OSOO】 Soonmelt® 375mg/Tab

ATC Code : J01CR02

中文名：雙合黴素膜衣錠 «永信»

適應症：葡萄球菌、鏈球菌、肺炎雙球菌、腦膜炎球菌及其他具有感受性細菌引起之感染症。

成分： Each tablet contains:
Amoxicillin 250 mg
Clavulanic Acid 125 mg

藥理分類： Antibiotic, Penicillin; Clavulanic acid: Beta-lactamases Inhibitor.

用法用量： **Note:** Dosage strengths based on amoxicillin component.

Treatment should not be extended beyond 14 days without review.

Administration:

Administer with food to increase absorption and decrease stomach upset.

Dosage regimens:

— Adults & Children $\geq$ 40 kg:

Mild to moderate infections:

500 mg every 12 hours or 250 mg every 8 hours.

Severe infections:

875 mg every 12 hours or 500 mg every 8 hours.

— Children < 40 kg:

- Soonmelt 375 毫克錠劑**不建議用於** 12 歲及 12 歲以下的兒童。(仿單)
- The 250-mg tablet of AUGMENTIN should not be used until the child weighs at least **40 kg**. (AUGMENTIN 原廠仿單)

Dose adjustment in renal impairment:

— Adults:

Note: Renally adjusted dose recommendations are based on the amoxicillin 250 mg/clavulanate 125 mg and amoxicillin 500 mg/clavulanate 125 mg tablets.

CrCl $\geq$ 30 mL/minute: No dosage adjustment necessary.

CrCl 10 to 30 mL/minute: 250 to 500 mg every 12 hours.

CrCl < 10 mL/minute: 250 to 500 mg every 24 hours.

End-stage renal disease (ESRD) on hemodialysis: 250 to 500 mg amoxicillin every 24 hours; administer dose both during and after dialysis.

— Children:

Similar reductions in dosage should be made for children.

不良反應：腹瀉(最常見)、消化不良、噁心、嘔吐、紅斑疹、蕁麻疹等。

注意事項：

1. Soonmelt 375 毫克錠劑**不建議用於** 12 歲及 12 歲以下的兒童。
2. Soonmelt® Tablet 易潮解，不宜磨粉。使用前再拆開包裝拿出藥錠。
3. 為減少胃腸不適之可能，可於**用餐開始時服用**，並可使藥劑吸收達到最大。

懷孕期：臨床判斷益處大於風險時才於懷孕時投與此藥。

授乳期：會有微量的藥物出現在乳汁，但除了會有感受度升高的危險之外，對以母乳餵哺的嬰兒並無任何不良影響。