Imidapril HCl

[OTAN] Tanatril® 5mg/Tab

中文名: 田納滋錠 《臺灣田邊》

適應症: 高血壓。

藥理分類: Angiotensin-Converting Enzyme (ACE) Inhibitor

用法用量: Administration: orally, taken with or without food.

Indications and dosage regimens:

Hypertension:

Initially, 5 mg once daily; maintenance, 10 mg once daily. May up to 20mg daiy.

ATC Code: C09AA16

Severe hypertension:

Initially 2.5 mg once daily.

不良反應: 頭痛、頭昏、疲倦、心悸、腹瀉、乾咳等。

交互作用: ● Potassium sparing diuretics, potassium supplements: ↑ in serum potassium.

• Lithium: 1 lithium concentration, potentially to toxic levels.

• NSAIDs: ↓ of the antihypertensive effect.

■ Insulin, hypoglycaemic sulphonamides: ↑ the hypoglycaemic effect.

• Tricyclic antidepressants, neuroleptics: ↑ antihypertensive effect and risk of orthostatic hypotension.

• Rifampicin: 1 antihypertensive effect.

Antacids: ↓ bioavailability of imidapril.

注意事項: 1.手術前 24 小時不建議投予。

2.嚴重腎功能受損者($CCr \le 30 \text{ mL/min}$,或血清中 creatinine 濃度 > 3 mg/dL), 須小心給藥,可考慮降低為正常劑量的一半或延長給藥間隔。

3.降壓作用可能導致頭暈或頭昏眼花,因此從事須有警覺性的活動(如操作機器 或駕車),需特別注意。

4.懷孕中期及末期投予 ACEIs,可能造成胎兒或新生兒的死亡或傷害,故禁用於孕婦及可能懷孕的婦女。

懷孕期: 1. 本品禁用於孕婦或可能已懷孕之婦人。

- 2. Exposure to an angiotensin-converting enzyme (ACE) inhibitor during pregnancy may be associated with an increased risk of fetal malformations.
- 3. Drugs that act on the renin-angiotensin system can cause injury and death to the developing fetus. Discontinue as soon as possible once pregnancy is detected.

授 乳 期: 動物(rat)試驗顯示 Tanatril 會移行入乳汁中,所以授乳婦女不應服用,不得已須 投與時應停止授乳。