

Imidapril HCl

【OTAN】Tanatril® 5mg/Tab

ATC Code : C09AA16

中文名：田納滋錠 «臺灣田邊»

適應症：高血壓。

藥理分類：Angiotensin-Converting Enzyme (ACE) Inhibitor

用法用量：**Administration:** orally, taken with or without food.

Indications and dosage regimens:

Hypertension:

Initially, 5 mg once daily; maintenance, 10 mg once daily. May up to 20mg daiy.

Severe hypertension:

Initially 2.5 mg once daily.

不良反應：頭痛、頭昏、疲倦、心悸、腹瀉、乾咳等。

- 交互作用：
- Potassium sparing diuretics, potassium supplements: ↑ in serum potassium.
 - Lithium: ↑ lithium concentration, potentially to toxic levels.
 - NSAIDs: **attenuation** of the antihypertensive effect.
 - Insulin, hypoglycaemic sulphonamides: ↑ the hypoglycaemic effect.
 - Tricyclic antidepressants, neuroleptics: ↑ antihypertensive effect and risk of orthostatic hypotension.
 - Rifampicin: **reduced** antihypertensive effect.
 - Antacids: **decreased** bioavailability of imidapril.

- 注意事項：
- 1.手術前 24 小時不建議投予。
 - 2.嚴重腎功能受損者($CCr \leq 30$ mL/min，或血清中 creatinine 濃度 > 3 mg/dL)，須小心給藥，可考慮降低為正常劑量的一半或延長給藥間隔。
 - 3.降壓作用可能導致頭暈或頭昏眼花，因此從事須有警覺性的活動（如操作機器或駕車），需特別注意。
 - 4.懷孕中期及末期投予 ACEIs，可能造成胎兒或新生兒的死亡或傷害，故禁用於孕婦及可能懷孕的婦女。

- 懷孕期：
- 1.本品禁用於孕婦或可能已懷孕之婦人。
 2. Exposure to an angiotensin-converting enzyme (ACE) inhibitor during pregnancy may be associated with an increased risk of fetal malformations.
 3. Drugs that act on the renin-angiotensin system can cause injury and death to the developing fetus. Discontinue as soon as possible once pregnancy is detected.

授乳期：動物(rat)試驗顯示 Tanatril 會移行入乳汁中，所以授乳婦女不應服用，不得已須投與時應停止授乳。