Iron (Hydroxide Polymaltose Complex)

【OTED】 Tedalin® Chewable Tab, 100mg/Tab

ATC Code : B03AB

中文名: 鐵達寧咀嚼錠 《元宙化學》

適應症: 預防及治療鐵質缺乏症、缺鐵性貧血症。

成 分: Each Tab contains:

Iron(III) (Hydroxide Polymaltose Complex)100 mg

藥理分類: Iron salt.

用法用量: Administration:

• Administer without regard to meals.

• Chew the tablet and then swallow with enough water.

Dosage regimen:

1~2 tab daily

不良反應: 噁心、皮膚起紅疹、嘔吐、腹瀉、心口疼痛、或臉部及四肢泛紅。

交互作用:

• Fluoroquinolones: ↓ effectiveness.

• Doxycycline, Tetracycline: ↓ tetracycline and iron effectiveness.

• Carbidopa or levodopa: ↓ carbidopa or levodopa effectiveness.

• Entacapone: ↓ entacapone bioavailability.

Proton-Pump Inhibitors: ↓ non-heme iron bioavailability.

注意事項: 1. 可空腹或與食物一起服用服用。

2. 咀嚼後和水一起吞服。

3. 服用後若糞便呈黑色屬正常現象,但若伴隨腹部劇烈疼痛,需立即就醫。

懷 孕 期: 1.Maternal iron requirements increase during pregnancy.

2. Ferrous gluconate has been evaluated in multiple studies as an iron supplement or for the treatment of IDA in pregnancy (*Peña-Rosas 2015; Reveiz 2011*).

授乳期: 1. Iron is present in breast milk (IOM 2001).

2. Maternal iron requirements are increased in breastfeeding women (IOM 2001).

3. The World Health Organization considers ferrous salts used for anemia to be compatible with breastfeeding (*WHO 2002*).