

Tonful (Acetaminophen, Carisoprodol)

【OPAN5】Tonful Tab[®]

ATC Code : M03BA52

中文名： 痛福錠 «永信藥品»

主成分： Each tablet contains:
Acetaminophen.....350mg
Carisoprodol.....175mg

適應症： 神經痛、關節痛、腰痛、背痛、扭傷、肌肉痛、脊椎炎、肌肉痙攣、關節周圍炎、斜頸。

藥理分類： **Analgesic, Nonopioid; Antipyretics; Skeletal Muscle Relaxant.**

作用機轉： 1.**Analgesic effects** are believed to be due to activation of descending serotonergic inhibitory pathways in the CNS. **Antipyresis** is produced from inhibition of the hypothalamic heat-regulating center.
2.Carisoprodol blocks interneuronal activity and depresses polysynaptic neuron transmission in the spinal cord and reticular formation of the brain.

用法用量： **Administration:** Orally, without regard to meals.

Dosage :

Adult:

1~2 tablets TID

不良反應： 黃疸、噁心、嘔吐、胃部痙攣或胃痛、上腹漲或痛。

注意事項： 1.兩次使用間隔請勿短於 4 小時。
2.習慣性飲酒之患者，應小心投與本藥，因 acetaminophen 可能造成肝傷害。
3.12 歲以下兒童不建議使用本藥。

懷 孕 期： 1. Acetaminophen crosses the placenta (Naga Rani 1989).
2. An association between maternal acetaminophen use and cryptorchidism (undescended testis) has been observed (Fisher 2016; Snijder 2012).
3. Acetaminophen is considered appropriate for the treatment of pain and fever in pregnancy (SMFM 2017).
4. Acetaminophen may be used as part of a multimodal approach to pain relief following cesarean delivery (ACOG 2019), for the treatment of acute migraine in pregnant patients (Burch 2019; Hamilton 2019a) and is recommended for the treatment of fever in pregnant women diagnosed with influenza (ACOG 753 2018).

授 乳 期： 1.Acetaminophen 會分泌至乳汁中，但 AAP 認為接受本藥治療期間之授乳婦可繼續哺乳。
2. Acetaminophen is one of the preferred non-narcotic agents (Sachs 2013) and is considered compatible with breastfeeding when used in usual recommended doses (WHO 2002).
3. Carisoprodol and its active metabolite, meprobamate, are present in breast milk.