Pentoxifylline

[OTRE1] Papiror® 100mg/Tab ATC Code : C04AD03

中文名: 排必樂腸溶錠 《杏林新生》

適應症: 末稍血管循環障礙。

[OPENT] Pentop® S.R. 400mg/Tab ATC Code: C04AD03

中文名: 循妥斯持續性錠 《優生》

適應症: 末稍血管循環障礙。

藥理分類: Blood Viscosity Reducer Agent.

用法用量: Administration:

Orally, preferably with meals. Swallow whole, do not crush or chew.

Indications and dosage regimens:

Intermittent claudication associated with peripheral vascular disease (i.e., chronic occlusive arterial disease of the extremities):

100~400mg TID.

- Beneficial effects may occur within 2-4 weeks; however, treatment should continue for at least 8 weeks to determine efficacy.
- Reduce dosage to 400 mg twice daily if adverse GI and / or CNS effects develop. Discontinue if adverse effects persist at this lower dosage. Efficacy was demonstrated in clinical studies of 6-months duration.

不良反應: 消化不良、噁心、嘔吐、腹瀉、眩暈、頭痛等。

交互作用:

Pentoxifylline may ↑ the antiplatelet effect of Agents with Antiplatelet Properties (Clopidogrel, Prasugrel, Aspirin, Dipyridamole, NSAIDs, Sulfinpyrazone, Ticagrelor, Ticlopidine, Tirofiban, Venlafaxine, Fluoxetine, Paroxetine, Citalopram, Sertraline, Piracetam, Cilostazol).

注意事項: 1、100 mg Trental 為腸溶劑型,宜整粒吞服,不可磨粉。

2、400 mg Trental 為緩釋劑型,宜整粒吞服,不可磨粉。

3、本品禁用於視網膜出血、腦出血、急性心肌梗塞及懷孕之患者。

4、腎功能不全之患者應減少劑量。

懷 孕 期: 孕婦不可服用本劑。

授 乳 期: 接受本藥治療期間,應停止授乳。