

Pentoxifylline

【OTRE1】Papiror® 100mg/Tab

ATC Code : C04AD03

中文名：排必樂腸溶錠 «杏林新生»

適應症：末稍血管循環障礙。

【OPENT】Pentop® S.R. 400mg/Tab

ATC Code : C04AD03

中文名：循妥斯持續性錠 «優生»

適應症：末稍血管循環障礙。

藥理分類：**Blood Viscosity Reducer Agent.**

用法用量：**Administration:**

- Orally, preferably with meals. Swallow whole, do not crush or chew.
- An aluminum and magnesium hydroxides antacid can be administered concomitantly to reduce intolerable GI effects.

Indications and dosage regimens:

Intermittent claudication associated with peripheral vascular disease (i.e., chronic occlusive arterial disease of the extremities):

100~400mg TID.

- Beneficial effects may occur within 2-4 weeks; however, treatment should continue for at least 8 weeks to determine efficacy.
- Reduce dosage to 400 mg twice daily if adverse GI and / or CNS effects develop. Discontinue if adverse effects persist at this lower dosage. Efficacy was demonstrated in clinical studies of 6-months duration.

不良反應：消化不良、噁心、嘔吐、腹瀉、眩暈、頭痛等。

交互作用：

- Pentoxifylline may ↑ the antiplatelet effect of Agents with Antiplatelet Properties (**Clopidogrel, Prasugrel, Acemetacin, Aspirin, Dipyridamole, Etodolac, Ibuprofen, Indomethacin, Ketoprofen, Ketorolac, Mefenamic Acid, Meloxicam, Naproxen, Piroxicam, Sulfipyrazone, Ticagrelor, Ticlopidine, Tirofiban, Venlafaxine, Fluoxetine, Paroxetine, Citalopram, Sertraline, Piracetam, Cilostazol**).

注意事項：1、100 mg Trental 為腸溶劑型，宜整粒吞服，不可磨粉。
2、400 mg Trental 為緩釋劑型，宜整粒吞服，不可磨粉。
3、本品禁用於視網膜出血、腦出血、急性心肌梗塞及懷孕之患者。
4、腎功能不全之患者應減少劑量。

懷孕期：孕婦不可服用本劑。

授乳期：接受本藥治療期間，應停止授乳。

使用規定：全民健保藥品給付規定—腦循環改善劑、腦代謝改善劑及血球流變改善劑：
因治療上需要使用腦循環改善劑或腦代謝改善劑或血球流變改善劑時，最多使用一類，且須檢附相關之紀錄或報告，並於病歷上詳細記載。