

Ritodrine HCl

【OYUT】Anpo® 10mg/Tab

ATC Code : G02CA01

中文名：安寶錠 «信東»

適應症：預防早產、流產。

藥理分類：Beta-2 agonist, Inhibition of acute preterm labor agent.

用法用量：**Administration** : Orally taken with or without food.

Indications and dosage regimens :

Maintenance

- May be started by administering the dose 30 mins before discontinuing the IV infusion.
- For the first 24 hrs of oral administration the usual dose is 10 mg ritodrine every 2 hrs.
- Thereafter, the usual maintenance dose is 10 to 20 mg every 4 to 6 hrs. The total daily oral dose of ritodrine should not exceed 120 mg.

不良反應：胸痛，呼吸困難，或視覺模糊，頭昏，口乾，皮膚乾、潮紅，心悸，頻尿，嘔吐，低血鉀。

注意事項：1.飯前或飯後服用均可。
2.用藥期間 須監測血壓及血鉀。

懷 孕 期：1.Ritodrine can cross the placenta.
2.There are no well-controlled studies using ritodrine before the 20th week of gestation, and the manufacturer recommends ritodrine not be used before the that time.

授 乳 期：1.Ritodrine is used to inhibit premature labor, and it is therefore unlikely that a lactating female would receive this drug.
2.There is insufficient evidence to support or refute its safety in breastfeeding. It is not known whether ritodrine is excreted into human breast milk and the potential for adverse effects in the nursing infant from exposure to the drug are unknown.