

Sevoflurane

【T1SOJ】 Sojourn Inhalation Liquid 250mL/Bot

ATC Code : N01AB08

中文名：舒眾吸入用液劑 «旌宇藥業»

適應症：吸入性全身麻醉。

藥理分類：**General Anesthetic, Inhalation.**

用法用量：**Anesthesia:**

Inhalation: Surgical levels of anesthesia are generally achieved with concentrations from 0.5% to 3% with or without the concomitant use of nitrous oxide; the concentration at which amnesia and loss of awareness occur is 0.6%.

MAC values for surgical levels of anesthesia:

Adult

25 years:

Sevoflurane in oxygen: 2.6%

Sevoflurane in 65% N2O/35% oxygen: 1.4%

40 years:

Sevoflurane in oxygen: 2.1%

Sevoflurane in 65% N2O/35% oxygen: 1.1%

60 years:

Sevoflurane in oxygen: 1.7%

Sevoflurane in 65% N2O/35% oxygen: 0.9%

80 years:

Sevoflurane in oxygen: 1.4%

Sevoflurane in 65% N2O/35% oxygen: 0.7%

Pediatric

MAC values for surgical levels of anesthesia:

0 to 1 month old full-term neonates: Sevoflurane in oxygen: 3.3%

1 to <6 months: Sevoflurane in oxygen: 3%

6 months to <1 year:

Sevoflurane in oxygen: 2.8%

Sevoflurane in 65% N2O/35% oxygen: 2%

1 to <3 years:

Sevoflurane in oxygen: 2.8%

Sevoflurane in 60% N2O/40% oxygen: 2%

3 to 12 years: Sevoflurane in oxygen: 2.5%

禁忌：已知對鹵素類麻醉劑過敏者禁用。已知或是懷疑易發生惡性高熱症者禁用。

不良反應：血壓降低、呼吸抑制；恢復期可能有顫抖、嘔吐、噁心。

注意事項：Isoflurane 會加強所有肌肉鬆弛劑的作用，尤其是對 nondepolarizing 類的肌肉鬆弛劑。當同時給予笑氣時，MAC 最小肺泡濃度會下降。

懷 孕 期：安全資料不足。除非明顯需要，否則不應於懷孕期間使用。

授 乳 期：目前尚不清楚本藥是否會在人乳中分泌。使用本藥 48 小時內不應哺乳。