

Atropine Sulfate

【TATRO1】Antol® 0.125% Eye Drops 5mL/Bot

ATC Code : S01FA01

中文名：安妥眼藥水 0.125% 《應元化學》

適應症：散瞳、睫狀肌麻痺。

藥理分類：Anticholinergic Agent, Ophthalmic; Ophthalmic Agent, Mydriatic.

用法用量：**Administration:**

- For topical ophthalmic use only.
- Wash hands before and after use. Avoid touching tip of applicator to eye or other surfaces.
- Finger pressure should be applied to lacrimal sac for 1 to 3 minutes after instillation to decrease risk of absorption and systemic reactions.

Indications and dosage regimens:

Mydriasis induction / Cycloplegic refraction:

- Adults: 1 drop of 0.25% ophthalmic solution in the eye 1 hour before examination.
- Children: 1-2 drops of 0.125% ophthalmic solution in the eye twice daily for 1-3 days before examination.

Uveitis(葡萄膜炎) / Iridocyclitis(虹膜睫狀體炎):

1-2 drops of 0.125% or 1% ophthalmic solution in the eye up to 3 times daily.

不良反應：眼睛刺痛、視力模糊、紅腫、癢、畏光、眼壓上升、口乾、心跳過快或頭痛。

注意事項：1.本劑禁用於青光眼之患者（對於狹角性青光眼的病人若使用 anticholinergic 及 adrenergic mydriatics 可能會導致急性的閉角性青光眼）。
2.使用本劑後會視力模糊，應避免從事駕駛或具危險性之機械操作。
3.開瓶 4 週後請勿繼續使用。

懷 孕 期：1.本品還未做動物生殖研究，也不知道是否能夠造成胎兒傷害。
2.本品僅當明確需要時才可應用於孕婦。
3. Atropine crosses the placenta following systemic maternal use (Shutt 1979).
4. Atropine is systemically available following ophthalmic administration.
5. If ophthalmic agents are needed during pregnancy, the minimum effective dose should be used in combination with punctual occlusion to decrease potential exposure to the fetus (Samples 1988).

授 乳 期：It is not known if atropine is excreted in breast milk following ophthalmic administration; however, systemic absorption occurs.

儲 存：請置於 25 °C 以下冷暗處避光儲存。