

Chloral Hydrate

【TCHL】Chloral Hydrate 10% Liquid 100mg/mL

ATC Code : N05CC01

藥理分類：Hypnotic, Miscellaneous.

用法用量：**Administration:** Rectal or PO

Indications and dosage regimens:

Sedation, Anxiety:

- Adults: 250 mg Q8h.
- Children: 5-15 mg/kg/dose Q8h. MAX 500 mg/dose.

Hypnotic:

- Adults: 500-1000 mg HS or 30 minutes prior to procedure, not to exceed 2 g/24hr.
- Children: 20-40 mg/kg/dose up to a MAX 50 mg/kg/24hr or 1 g/dose or 2 g/24hr.

Prior to EEG:

- Children: 20-25 mg/kg/dose, 30-60 minutes prior to EEG; may repeat to MAX 100 mg/kg or 2 g total.

Sedative, nonpainful procedure: PO

- Children: 50-75 mg/kg/dose 30-60 min prior to procedure; may repeat 30 min after initial dose if need, to a total MAX 120 mg/kg.

不良反應： 心律不整、睡眠形態改變、呼吸抑制、皮膚過敏。

注意事項： 可能引起頭暈或嗜睡，開車或操作危險機械時藥小心。

懷 孕 期： Animal reproduction studies have not been conducted. Chloral hydrate crosses the placenta, and long-term use may lead to withdrawal symptoms in the neonate.

授 乳 期： 1.Chloral hydrate is excreted in breast milk; use by breast-feeding women may cause sedation in the infant.
2.謹慎使用。

備 註： 本品以原料 chloral hydrate 配製。