Ketoconazole

[TKET] Ketoconazole® Cream 20mg/g, 10g/Tube ATC Code: D01AC08

中文名: 必克多黴乳膏 《杏辉》

適應症: 皮膚黴菌和酵母菌所引起的皮膚、頭髮及指甲感染症。

藥理分類: Antifungal Agent, Imidazole Derivative; Topical.

用法用量: Apply once daily to the affected areas of the skin.

Therapy should be continued for 2 weeks in order to minimize the risk of recurrence. Safety and efficacy of 2% cream not established in children; 2% cream has been used

without unusual adverse effect in children 2 days to 12 years of age.

不良反應: 皮膚局部反應(刺激感、灼熱感)。

注意事項: 若同時使用類固醇藥膏,可於白天使用類固醇藥膏,晚上使用 Ketoconazole

Cream.

懷 孕 期: 1.局部使用不會經由皮膚吸收,可用於孕婦(仿單)。

2. Topical ketoconazole may be used for the treatment of infectious atopic dermatitis

complications in pregnant women (Vestergaard 2019).

授 乳 期: 1.局部使用不會經由皮膚吸收,可用於哺乳婦(仿單)。

2. Do not apply to the nipple or surrounding area to avoid direct contact with infants

who are breastfeeding.