

Mometasone Furoate

【TMETC】Metsone[®] Cream 0.1% 15g/Tube

ATC Code : D07AC13

中 文 名 : "杏輝" 頓安膚乳膏 «杏輝»

主 成 分 : Each gram contain:
Mometasone Furoate 1mg

適 應 症 : 對類固醇具有感受性之皮膚炎、皮膚癢。

藥理分類: **Corticosteroid, Topical.**

用法用量: A thin film of Metsone[®] Cream 0.1% should be applied to the affected skin areas once daily until the lesion heals or for a duration of 3 weeks.

不良反應: 局部感覺異常、癢、皮膚萎縮。

注意事項: 懷孕婦女需小心使用。

懷 孕 期: 1. 頓安膚對懷孕婦女的安全性尚未建立，因此只有在對母體的有益性大於對胎兒的危險性時才可使用局部類固醇。此等級的藥品不得大量或長期使用於孕婦。
2. Adverse events have been observed in animal reproduction studies.
3. Use of mild to moderate potency topical corticosteroids is preferred in pregnant females and the use of large amounts or use for prolonged periods of time should be avoided (Chi 2016; Chi 2017; Murase 2014).
4. Also avoid areas of high percutaneous absorption (Chi 2017).

授 乳 期: 1. 局部使用類固醇而全身吸收，尚不知是否會分泌在乳汁中；全身性的使用類固醇會分泌於乳汁中。
2. It is not known if sufficient quantities of mometasone are absorbed following topical administration to produce detectable amounts in breast milk.
3. Topical corticosteroids are generally considered acceptable for use in breastfeeding females (Butler 2014; WHO 2002).
4. Do not apply topical corticosteroids to breast until breastfeeding ceases (Leachman 2006);