

Mycomb (Triamcinolone, Neomycin, Gramicidin, Nystatin)

【TTAR】Mycomb® Otic Drops, 5mL/Bot

ATC Code : S02CA04

中文名：杏輝美康耳用滴劑 «杏輝»

適應症：外耳道念珠菌感染、表面細菌感染、異位性皮膚炎、濕疹性皮膚炎、過敏接觸性皮膚炎、原發刺激性接觸皮膚炎、皮脂溢出性皮膚炎、溢出性皮膚炎、神經性皮膚炎。

成分：Each mL contains:

Triamcinolone Acetonide.....1mg
Neomycin (as Sulfate) 2.5mg
Gramicidin.....0.25mg
Nystatin.....100,000units

藥理分類：Antibiotic, Topical; Antibiotic/Corticosteroid, Topical; Antifungal Agent, Topical; Corticosteroid, Topical. 抗炎、止癢殺菌和殺黴菌。

用法用量：Administration:

- 用於治療耳部及其周圍皮膚各。

Indications and Dosage regimens:

每天使用 3~4 次，每次 2~3 滴，滴入耳道，或以棉球沾之塞於耳道內使用。

不良反應：耳刺激感、搔癢。頭暈。

注意事項：

1. 對本產品所含成分過敏之病人及鼓膜穿孔之病人(因含有 aminoglycoside 類成分)，不建議使用。
2. 如果藥水冰冷，滴入時容易發生暈眩的情形，故儘可能在與體溫相近的情形下使用本劑，滴入後讓藥水停留約 10 分鐘。
3. 本品不得用於皮膚結核，急性單純疱疹，天花，水痘，亦不可用以點眼及眼圈附近塗敷。

懷 孕 期：1. Animal reproduction studies have not been conducted with this combination.
2. Aminoglycosides cross the placenta. Aminoglycosides may cause fetal harm if administered to a pregnant woman.
3. Absorption following oral use is poor and nystatin is not absorbed following application to mucous membranes or intact skin.

授 乳 期：1. It is not known if neomycin is excreted into breast milk.
2. Absorption following oral use is poor and nystatin is not absorbed following application to mucous membranes or intact skin.
3. The manufacturer recommends that caution be exercised when administering nystatin to nursing women.

儲 存：於 25°C 以下室溫保存。