Mometasone Furoate

[TNAS] Nasonex® Aqueous Nasal spray 50mcg/Spray

ATC Code: R01AD09

中文名: 內舒拿水溶性鼻用噴液劑 《先靈葆雅》

適應症: 成人青少年及二歲以上兒童之季節性或常年性過敏性鼻炎。亦用於治療 18 歲及

以上成人輕度到中度鼻息肉之相關症狀。

藥理分類: Corticosteroid, Nasal.

用法用量: **Administration:** For intranasal administration only.

- Prime pump (press 10 times or until fine spray appears) prior to first use. If 7 or more days have elapsed since last use, reprime pump with 2 sprays or until a fine mist appears.
- Shake before using. Blow nose to clear nostrils. Insert applicator into nostril, tilt head slightly forward keeping bottle upright, and close off the other nostril. Breathe in through nose.
- While inhaling, press pump to release spray; exhale through mouth.
- After each use, wipe the spray tip with a clean tissue and replace cap.

Indications and dosage regimens:

Seasonal and perennial allergic rhinitis: Intranasal:

Adult & children ≥ 12 years:

Treatment: 2 sprays (100 mcg) in each nostril once daily.

Prevention: 2 sprays (100 mcg) in each nostril once daily beginning 2-4 weeks prior to pollen season.

Children 2-11 years: 1 spray (50 mcg) in each nostril once daily.

Treatment of nasal polyps: Intranasal: ≥18 years:

2 sprays (100 mcg) in each nostril twice daily; 2 sprays (100 mcg) once daily may be effective in some patients.

不良反應: 頭痛、鼻腔出血、咽喉炎、鼻腔刺激。

注意事項: 1.本品僅供噴鼻使用,使用前請先搖勻。

懷 孕 期: 懷孕婦女,只有在評估過使用的利益勝於對母親、胎兒的危險性後,才可以給

予本品。

授 乳 期: 授乳婦女只有在評估過使用的利益勝於對母親、新生兒的危險性後,才可以給

予本品。